

MATH 140 EXAM #1 KEY (SPRING 2022)

1 We have

$$\lim_{x \rightarrow -1} F(x) = 3, \quad \lim_{x \rightarrow -2} F(x) = 1, \quad \lim_{x \rightarrow 1^-} F(x) = 2, \quad \lim_{x \rightarrow 1^+} F(x) = 4, \quad \lim_{x \rightarrow 3^+} F(x) = \text{DNE}.$$

2a Simply reduce the fraction first:

$$\lim_{x \rightarrow -b} \frac{(x+b)^6 + (x+b)^9}{4} = 0.$$

2b Multiply top and bottom of fraction by $5(5+h)$:

$$\lim_{h \rightarrow 0} \frac{5 - (5+h)}{5h(5+h)} = \lim_{h \rightarrow 0} \frac{-1}{5(5+h)} = -\frac{1}{25}.$$

2c Factor the bottom (or multiply top and bottom by $\sqrt{x}+9$):

$$\lim_{x \rightarrow 81} \frac{\sqrt{x}-9}{(\sqrt{x}-9)(\sqrt{x}+9)} = \lim_{x \rightarrow 81} \frac{1}{\sqrt{x}+9} = \frac{1}{\sqrt{81}+9} = \frac{1}{18}.$$

2d Factor the bottom:

$$\lim_{x \rightarrow 0} \frac{1 - \cos x}{(\cos x - 2)(\cos x - 1)} = \lim_{x \rightarrow 0} \frac{-1}{\cos x - 2} = \frac{-1}{\cos 0 - 2} = 1.$$

3 Since

$$\lim_{x \rightarrow -3^-} G(x) = \lim_{x \rightarrow -3^-} (3x + 2k) = -9 + 2k \quad \text{and} \quad \lim_{x \rightarrow -3^+} G(x) = \lim_{x \rightarrow -3^+} (x + 9) = 6,$$

the limit $\lim_{x \rightarrow -3} G(x)$ can only exist if $-9 + 2k = 6$, which only happens if $k = \frac{15}{2}$. Then $\lim_{x \rightarrow -3} G(x) = 6$.

4 All equal $-\infty$.

5 Since

$$f(x) = \frac{x+1}{x(x-2)^2},$$

f has vertical asymptotes $x = 0$ and $x = 2$. We find that

$$\lim_{x \rightarrow 0^-} f(x) = -\infty, \quad \lim_{x \rightarrow 0^+} f(x) = +\infty, \quad \lim_{x \rightarrow 2^-} f(x) = +\infty, \quad \lim_{x \rightarrow 2^+} f(x) = +\infty,$$

so $\lim_{x \rightarrow 2} f(x) = +\infty$ and $\lim_{x \rightarrow 0} f(x)$ does not exist.

6 Divide top and bottom by x^2 , then evaluate to get $-\frac{4}{7}$.

7a In general,

$$f(x) = \frac{4x^3 + 1}{2x^3 + \sqrt{16x^6 + 1}} = \frac{4x^3 + 1}{2x^3 + |x|^3\sqrt{16 + x^{-6}}},$$

so

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} \frac{4x^3 + 1}{2x^3 + x^3\sqrt{16 + x^{-6}}} = \lim_{x \rightarrow \infty} \frac{4 + x^{-3}}{2 + \sqrt{16 + x^{-6}}} = \frac{4}{2 + \sqrt{16}} = \frac{2}{3},$$

and

$$\lim_{x \rightarrow -\infty} f(x) = \lim_{x \rightarrow -\infty} \frac{4x^3 + 1}{2x^3 - x^3\sqrt{16 + x^{-6}}} = \lim_{x \rightarrow -\infty} \frac{4 + x^{-3}}{2 - \sqrt{16 + x^{-6}}} = \frac{4}{2 - \sqrt{16}} = -2.$$

7b Horizontal asymptotes are $y = \frac{2}{3}$ and $y = -2$.

8 Continuity from the left at -1 requires that $\lim_{x \rightarrow -1^-} h(x) = h(-1)$. Since

$$\lim_{x \rightarrow -1^-} h(x) = \lim_{x \rightarrow -1^-} (2x^2 + x) = 1,$$

and $h(-1) = s$, we set $s = 1$ to get continuity from the left at -1 .

Continuity from the right at -1 requires $\lim_{x \rightarrow -1^+} h(x) = h(-1)$. Since

$$\lim_{x \rightarrow -1^+} h(x) = \lim_{x \rightarrow -1^+} (4 - 3x) = 7,$$

and $h(-1) = s$, we set $s = 7$ to get continuity from the right at -1 .

9a By definition,

$$f'(4) = \lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4} = \lim_{x \rightarrow 4} \frac{\sqrt{x-3} - 1}{x - 4} = \lim_{x \rightarrow 4} \frac{x - 4}{(x - 4)(\sqrt{x-3} + 1)} = \lim_{x \rightarrow 4} \frac{1}{\sqrt{x-3} + 1} = \frac{1}{2}.$$

9b Slope of the tangent line at $(4, 1)$ is $f'(4) = \frac{1}{2}$, so line is $y = \frac{1}{2}x - 1$.