

1. 10 pts. each Use integration by parts (possibly more than once) to find each integral.

(a) $\int x \cos 8x \, dx$

(b) $\int e^{-x} \sin 4x \, dx$

(c) $\int_{2/\sqrt{3}}^2 z \sec^{-1}(z) \, dz$

2. 10 pts. each Determine each indefinite trigonometric integral.

(a) $\int \sin^7(x) \cos^3(x) \, dx$

(b) $\int \frac{\csc^4 t}{\cot^2 t} \, dt$

(c) $\int_{-\pi/3}^{\pi/3} \sqrt{\sec^2(\varphi) - 1} \, d\varphi$

3. 10 pts. each Use a trigonometric substitution to determine the indefinite integral

(a) $\int_0^{1/3} \frac{1}{(9x^2 + 1)^{3/2}} \, dx$

(b) $\int \sqrt{121 - x^2} \, dx$

4. 10 pts. each Use partial fractions to determine the indefinite integral

(a) $\int \frac{y + 1}{y^3 + 3y^2 - 18y} \, dy$

(b) $\int \frac{2}{(x - 4)(x^2 + 2x + 6)} \, dx$

5. 10 pts. each Evaluate the improper integral or state that it diverges.

(a) $\int_{-\infty}^1 \frac{1}{(2x - 3)^2} \, dx$

(b) $\int_0^4 \frac{1}{x - 1} \, dx$

6. 10 pts. Let $\mathcal{R}$ be the region on the xy -plane bounded by $f(x) = x^{-2}$ and the x -axis on the interval $[2, \infty)$. Find the volume of the solid generated by revolving $\mathcal{R}$ about the x -axis.

FORMULAS & DEFINITIONS

1. $\theta = \tan^{-1} x \Leftrightarrow x = \tan \theta$, for $\theta \in (-\pi/2, \pi/2)$
2. $\theta = \cot^{-1} x \Leftrightarrow x = \cot \theta$, for $\theta \in (0, \pi)$
3. $\theta = \sec^{-1} x \Leftrightarrow x = \sec \theta$, for $\theta \in [0, \pi/2) \cup (\pi/2, \pi]$
4. $\theta = \csc^{-1} x \Leftrightarrow x = \csc \theta$, for $\theta \in [-\pi/2, 0) \cup (0, \pi/2]$
5. $(b^x)' = b^x \ln b$, for $b \in (0, 1) \cup (1, \infty)$
6. $(\log_b |x|)' = \frac{1}{x \ln b}$, for $x \neq 0$
7. $(\sin^{-1} x)' = \frac{1}{\sqrt{1-x^2}}$, for $x \in (-1, 1)$
8. $(\tan^{-1} x)' = \frac{1}{1+x^2}$, for $x \in (-\infty, \infty)$
9. $(\sec^{-1} x)' = \frac{1}{|x|\sqrt{x^2-1}}$, for $x \in (-\infty, -1) \cup (1, \infty)$
10. $\int b^x dx = \frac{1}{\ln b} b^x + c$, for $b \in (0, 1) \cup (1, \infty)$
11. $\int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1} \left(\frac{x}{a} \right) + c$, for $a \in (0, \infty)$
12. $\int \frac{1}{a^2+x^2} dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + c$, for $a \neq 0$
13. $\int \frac{1}{x\sqrt{x^2-a^2}} dx = \frac{1}{a} \sec^{-1} \left| \frac{x}{a} \right| + c$, for $a \in (0, \infty)$
14. $\int \sin^n x dx = -\frac{\sin^{n-1} x \cos x}{n} + \frac{n-1}{n} \int \sin^{n-2} x dx$
15. $\int \cos^n x dx = \frac{\cos^{n-1} x \sin x}{n} + \frac{n-1}{n} \int \cos^{n-2} x dx$
16. $\int \tan^n x dx = \frac{\tan^{n-1} x}{n-1} - \int \tan^{n-2} x dx$, $n \neq 1$
17. $\int \sec^n x dx = \frac{\sec^{n-2} x \tan x}{n-1} + \frac{n-2}{n-1} \int \sec^{n-2} x dx$, $n \neq 1$
18. $\int \tan x dx = -\ln |\cos x| + c = \ln |\sec x| + c$
19. $\int \cot x dx = \ln |\sin x| + c$
20. $\int \sec x dx = \ln |\sec x + \tan x| + c$
21. $\int \csc x dx = -\ln |\csc x + \cot x| + c$