

1 We have

$$\begin{aligned}\mathcal{V} &= \int_{-2}^2 \int_{-\sqrt{1-x^2/4}}^{\sqrt{1-x^2/4}} \int_{x-3}^{3-x} dz dy dx = \int_{-2}^2 \int_{-\sqrt{1-x^2/4}}^{\sqrt{1-x^2/4}} (6-2x) dy dx \\ &= \int_{-2}^2 2(6-2x)\sqrt{1+x^2/4} dx = 2 \int_{-2}^2 (3-x)\sqrt{4-x^2} dx.\end{aligned}$$

Trigonometric substitution: let $x = 2 \sin \theta$, so

$$\begin{aligned}\mathcal{V} &= 2 \int_{-\pi/2}^{\pi/2} (3-2\sin \theta) \sqrt{4-4\sin^2 \theta} \cdot 2 \cos \theta d\theta = 24 \int_{-\pi/2}^{\pi/2} \cos^2 \theta d\theta - 16 \int_{-\pi/2}^{\pi/2} \sin \theta \cos^2 \theta d\theta \\ &= 12 \int_{-\pi/2}^{\pi/2} (1+\cos 2\theta) d\theta = 12 \left[\theta + \frac{1}{2} \sin 2\theta \right]_{-\pi/2}^{\pi/2} = 12\pi.\end{aligned}$$

2 Letting I be the integral, we have

$$\begin{aligned}I &= \int_1^6 \int_0^{4-2y/3} \left[\frac{x}{y} \right]_0^{12-2y-3z} dz dy = \int_1^6 \int_0^{4-2y/3} \frac{12-2y-3z}{y} dz dy \\ &= \int_1^6 \left[\frac{12z}{y} - 2z - \frac{3z^2}{2y} \right]_0^{4-2y/3} dy = \int_1^6 \left(\frac{24}{y} + \frac{2y}{3} - 8 \right) dy = 24 \ln 6 - \frac{85}{3}\end{aligned}$$

3 The region D is shown at left in the figure below. It will be convenient to work in cylindrical coordinates, where $x = r \cos \theta$ and $y = r \sin \theta$ so that the equation of the paraboloid becomes

$$z = x^2 + y^2 = (r \cos \theta)^2 + (r \sin \theta)^2 = r^2,$$

and the equation of the plane remains $z = 25$.

The intersection of the surfaces $z = 25$ and $z = x^2 + y^2$ is the set of points

$$\{(x, y, 25) : x^2 + y^2 = 25\},$$

which is a curve that projects onto the xy -plane as a circle of radius 5 centered at the origin. Thus, the projection of D onto the xy -plane is a region R that is a closed disc with radius 5 centered at the origin, shown at right in the figure below.

Now, a point in R may have a θ -coordinate value ranging anywhere from $\theta = 0$ to $\theta = 2\pi$; that is, if $(r, \theta) \in R$, then $0 \leq \theta \leq 2\pi$.

If we fix $\theta \in [0, 2\pi]$, then a point $(r, \theta) \in R$ must lie on the line segment joining $o = (0, 0)$ and $a = (5, \theta)$, shown at right in the figure below. That is, given $\theta \in [0, 2\pi]$, a point $(r, \theta) \in R$ can have r -coordinate value ranging anywhere from $r = 0$ to $r = 5$, which is to say $0 \leq r \leq 5$.

Finally, fixing $\theta \in [0, 2\pi]$ and $r \in [0, 5]$, we consider the limits on z in order for (r, θ, z) to be a point that lies in D . We find that generally z must be such that (r, θ, z) is above the paraboloid $z = r^2$ and below the plane $z = 25$, which is to say $r^2 \leq z \leq 25$.


Thus we find that the region $E \subseteq \mathbb{R}_{r\theta z}^3$ for which $T_{\text{cyl}}(E) = D$ is

$$E = \{(r, \theta, z) : 0 \leq \theta \leq 2\pi, 0 \leq r \leq 5, r^2 \leq z \leq 25\},$$

and so

$$\begin{aligned}
 \mathcal{V}(D) &= \iiint_D dV = \iiint_E r dV = \int_0^{2\pi} \int_0^5 \int_{r^2}^{25} r dz dr d\theta \\
 &= \int_0^{2\pi} \int_0^5 (25r - r^3) dr d\theta = \int_0^{2\pi} \left[\frac{25}{2} r^2 - \frac{1}{4} r^4 \right]_0^5 d\theta \\
 &= \int_0^{2\pi} \frac{625}{4} d\theta = \frac{625}{4} \cdot 2\pi = \frac{625}{2} \pi
 \end{aligned}$$

is the volume of the region D .



4 D is given as $\{(\rho, \varphi, \theta) : 0 \leq \theta \leq 2\pi, 0 \leq \varphi \leq \pi, 0 \leq \rho \leq 1\}$ in spherical coordinates, so

$$\iiint_D e^{-(x^2+y^2+z^2)^{3/2}} dV = \int_0^{2\pi} \int_0^\pi \int_0^1 e^{-\rho^3} \rho^2 \sin \varphi d\rho d\varphi d\theta.$$

Substitution: let $u = -\rho^3$, so that $\rho^2 d\rho = -\frac{1}{3} du$ and we obtain

$$\begin{aligned}
 \iiint_D e^{-(x^2+y^2+z^2)^{3/2}} dV &= \int_0^{2\pi} \int_0^\pi -\frac{\sin \varphi}{3} \left(\int_0^{-1} e^u du \right) d\varphi d\theta = \frac{1 - e^{-1}}{3} \int_0^{2\pi} \int_0^\pi \sin \varphi d\varphi d\theta \\
 &= \frac{1 - e^{-1}}{3} \int_0^{2\pi} 2 d\theta = \frac{1 - e^{-1}}{3} \cdot 4\pi = \frac{4\pi(e - 1)}{3e}
 \end{aligned}$$

5 The line may be parameterized by $\mathbf{r}(t) = \langle t, t \rangle$, $-\infty < t < \infty$, with tangent vector at $\mathbf{r}(t)$ given by $\mathbf{r}'(t) = \langle 1, 1 \rangle$. We must find a vector field \mathbf{F} such that, for each t , $\mathbf{F}(\mathbf{r}(t))$ is orthogonal (i.e. normal) to $\mathbf{r}'(t)$. That is, we must have

$$\mathbf{F}(t, t) \cdot \langle 1, 1 \rangle = 0.$$

There are many possibilities: $\mathbf{F}(x, y) = \langle -1, 1 \rangle$, or $\mathbf{F}(x, y) = \langle -c, c \rangle$ for any $c \neq 0$, or $\mathbf{F}(x, y) = \langle -x, x \rangle$. (A trivial solution would be $\mathbf{F}(x, y) = \langle 0, 0 \rangle$, but we are looking for a nonzero vector field here.)

6 Since $|\mathbf{r}'(t)| = \sqrt{10}$, we have

$$\int_C (y - z) = \int_0^{2\pi} (3 \sin t - t) \sqrt{10} dt = -2\pi^2 \sqrt{10}.$$

7 Here $\mathbf{r}'(t) = \langle -4 \sin t, 4 \cos t \rangle$, so

$$\begin{aligned} \int_C \mathbf{F} \cdot d\mathbf{r} &= \int_0^\pi \mathbf{F}(4 \cos t, 4 \sin t) \cdot \langle -4 \sin t, 4 \cos t \rangle dt \\ &= \int_0^\pi \langle -4 \sin t, 4 \cos t \rangle \cdot \langle -4 \sin t, 4 \cos t \rangle dt \\ &= \int_0^\pi 16 dt = 16\pi. \end{aligned}$$

8 For brevity let $\mathbf{x} = \langle x, y, z \rangle$. We have $\mathbf{F} = \langle f, g, h \rangle$ with $f(\mathbf{x}) = 2xy^3z^4$, $g(\mathbf{x}) = 3x^2y^2z^4$, and $h(\mathbf{x}) = 4x^2y^3z^3$. Since

$$f_y(\mathbf{x}) = 6xy^2z^4 = g_x(\mathbf{x}), \quad f_z(\mathbf{x}) = 8xy^3z^3 = h_x(\mathbf{x}), \quad g_z(\mathbf{x}) = 12x^2y^2z^3 = h_y(\mathbf{x}),$$

it follows that \mathbf{F} is conservative.

We now find φ . From $\nabla\varphi = \mathbf{F}$ we have

$$\varphi_x(\mathbf{x}) = 2xy^3z^4, \quad \varphi_y(\mathbf{x}) = 3x^2y^2z^4, \quad \varphi_z(\mathbf{x}) = 4x^2y^3z^3.$$

Hence

$$\varphi(\mathbf{x}) = \int \varphi_x(\mathbf{x}) dx = x^2y^3z^4 + c(y, z).$$

Differentiating this with respect to y gives $\varphi_y(\mathbf{x}) = 3x^2y^2z^4 + c_y(y, z)$, which, when compared to $\varphi_y(\mathbf{x}) = 3x^2y^2z^4$, informs us that $c_y(y, z) = 0$ and therefore $c(y, z) = c(z)$ (that is, the function c must not be a function of y).

At this point we have $\varphi(\mathbf{x}) = x^2y^3z^4 + c(z)$. This implies that $\varphi_z(\mathbf{x}) = 4x^2y^3z^3 + c'(z)$, which, when compared to $\varphi_z(\mathbf{x}) = 4x^2y^3z^3$, informs us that $c'(z) = 0$. So $c(z) = c$, where c is an arbitrary constant. Choosing c be zero, we obtain $\varphi(\mathbf{x}) = x^2y^3z^4$.

9 The curve C goes from $\mathbf{a} = \langle 0, 0 \rangle$ to $\mathbf{b} = \langle \ln 2, 2\pi \rangle$, and the fact that it's a line segment will be irrelevant. Letting $\varphi(x, y) = e^{-x} \cos y$, the Fundamental Theorem of Line Integrals gives

$$\begin{aligned} \int_C \nabla(e^{-x} \cos y) \cdot d\mathbf{r} &= \int_C \nabla\varphi \cdot d\mathbf{r} = \varphi(\mathbf{b}) - \varphi(\mathbf{a}) = \varphi(\ln 2, 2\pi) - \varphi(0, 0) \\ &= e^{-\ln 2} \cos(2\pi) - e^{-0} \cos(0) = \frac{1}{2} - 1 = -\frac{1}{2}. \end{aligned}$$